# Factoring by Trial and Error Method

—— Factor each trinomial using the trial and error method.  $\,-\,$ 

### Solved Example: $6x^2 - 25x + 24$

- Paired factors of  $6x^2$  are x(6x) or 2x(3x)
- Thus, the factors can be  $(x \underline{\hspace{1cm}})$   $(6x \underline{\hspace{1cm}})$  or  $(2x \underline{\hspace{1cm}})$   $(3x \underline{\hspace{1cm}})$
- Try all paired factors of 24
  Paired factors are (1 and 24, 2 and 12, 3 and 8, 4 and 6)
- So, the correct factors are (2x 3)(3x 8)

4x <sup>2</sup> - 5x - 6	$5x^2 + 8x - 4$
5x <sup>2</sup> - 22x - 15	$x^2 - 9x + 18$
$2x^2 + 3x - 2$	$x^2 - 8x + 16$

## Factoring by Trial and Error Method

#### **Answers**

### Solved Example: $6x^2 - 25x + 24$

- Paired factors of  $6x^2$  are x(6x) or 2x(3x)
- Thus, the factors can be  $(x \underline{\hspace{1cm}})$   $(6x \underline{\hspace{1cm}})$  or  $(2x \underline{\hspace{1cm}})$   $(3x \underline{\hspace{1cm}})$
- Try all paired factors of 24
  Paired factors are (1 and 24, 2 and 12, 3 and 8, 4 and 6)
- So, the correct factors are (2x 3)(3x 8)

4x <sup>2</sup> - 5x - 6	$5x^2 + 8x - 4$
(4x + 3)(x - 2)	(5x - 2)(x + 2)
5x <sup>2</sup> - 22x - 15	$x^2 - 9x + 18$
(5x + 3)(x - 5)	(x - 3)(x - 6)
$2x^2 + 3x - 2$	$x^2 - 8x + 16$
(2x - 1)(x + 2)	$(x - 4)^2$